

AQA Combined Science: **Trilogy**

Year 10

Knowledge organiser booklet

Paper 1

CHEMISTRY

C1 Atomic structure and the periodic table

C2 Bonding, structure and the properties of matter

C3 Quantitative chemistry

C4 Chemical changes

C5 Energy changes

Name:

The Periodic Table of Elements

1		2												3		4	5	6	7	0							
																		1 H hydrogen 1									4 He helium 2
7 Li lithium 3		9 Be beryllium 4		Key relative atomic mass atomic symbol name atomic (proton) number										11 B boron 5		12 C carbon 6	14 N nitrogen 7	16 O oxygen 8	19 F fluorine 9	20 Ne neon 10							
23 Na sodium 11		24 Mg magnesium 12												27 Al aluminium 13		28 Si silicon 14	31 P phosphorus 15	32 S sulfur 16	35.5 Cl chlorine 17	40 Ar argon 18							
39 K potassium 19		40 Ca calcium 20	45 Sc scandium 21	48 Ti titanium 22	51 V vanadium 23	52 Cr chromium 24	55 Mn manganese 25	56 Fe iron 26	59 Co cobalt 27	59 Ni nickel 28	63.5 Cu copper 29	65 Zn zinc 30	70 Ga gallium 31	73 Ge germanium 32	75 As arsenic 33	79 Se selenium 34	80 Br bromine 35	84 Kr krypton 36									
85 Rb rubidium 37		88 Sr strontium 38	89 Y yttrium 39	91 Zr zirconium 40	93 Nb niobium 41	96 Mo molybdenum 42	[98] Tc technetium 43	101 Ru ruthenium 44	103 Rh rhodium 45	106 Pd palladium 46	108 Ag silver 47	112 Cd cadmium 48	115 In indium 49	119 Sn tin 50	122 Sb antimony 51	128 Te tellurium 52	127 I iodine 53	131 Xe xenon 54									
133 Cs caesium 55		137 Ba barium 56	139 La* lanthanum 57	178 Hf hafnium 72	181 Ta tantalum 73	184 W tungsten 74	186 Re rhenium 75	190 Os osmium 76	192 Ir iridium 77	195 Pt platinum 78	197 Au gold 79	201 Hg mercury 80	204 Tl thallium 81	207 Pb lead 82	209 Bi bismuth 83	[209] Po polonium 84	[210] At astatine 85	[222] Rn radon 86									
[223] Fr francium 87		[226] Ra radium 88	[227] Ac* actinium 89	[261] Rf rutherfordium 104	[262] Db dubnium 105	[266] Sg seaborgium 106	[264] Bh bohrium 107	[277] Hs hassium 108	[268] Mt meitnerium 109	[271] Ds darmstadtium 110	[272] Rg roentgenium 111	Elements with atomic numbers 112 – 116 have been reported but not fully authenticated															


* The Lanthanides (atomic numbers 58 – 71) and the Actinides (atomic numbers 90 – 103) have been omitted.

Relative atomic masses for Cu and Cl have not been rounded to the nearest whole number.

C1 – Atomic Structure and The Periodic Table

Atoms

- Made up of **protons, electrons** and **neutrons**.




Subatomic particle	Relative Mass	Charge
Proton	1	Positive
Neutron	1	Neutral
Electron	Very small	Negative

Atoms have a radius of about 0.1nm (1×10^{-10} m)

Radius of nucleus = about 1×10^{-14} m

Elements

- Only have **one type** of atom
- Found on the **Periodic Table**




Compounds

- Two or more elements **chemically combined**.
- Formed by chemical reactions
- For example: CO_2 H_2O CH_4 HCl NaCl

Isotopes

Isotope = atoms of the **same element** which have the **same number of protons**, but a **different number of neutrons**.

These are isotopes because..



Chemical Equations

- Shown by using a **word equation**.
e.g. magnesium + oxygen $\rightarrow$ magnesium oxide

Left of the arrow = **reactants**
Right of the arrow = **products**.

- Also can be shown by a **symbol equation**
e.g. $2\text{Mg} + \text{O}_2 \rightarrow 2\text{MgO}$

Mixtures and Separation


Mixtures – two or more elements or compounds **not** chemically joined.

This means the different components of the mixture can be separated by physical methods (below)

E.g. air is a mixture mainly made of nitrogen, oxygen and carbon dioxide.

Chromatography

to separate out mixtures (usually liquids) (e.g. colours in ink)



Filtration

To separate insoluble solids from liquids (e.g. sand and water)



Evaporation

To quickly separate soluble solids from a solution. (e.g. salt and water)



Crystallisation


To slowly separate a soluble salt from a solution. (e.g. copper sulfate crystals)



C1 – Atomic Structure and The Periodic Table

Distillation

Simple distillation – separating a liquid from a solution.




- Liquid is heated to boiling point and evaporates
- Vapours travel up into the condenser
- Condenser has cold water around it.
- Vapours cool and condense (turn back into a liquid).

Electronic Structure

- Electrons are found on shells (orbits) orbiting the nucleus.
- There is a maximum number of electrons allowed on each shell:


First shell = 2 electrons
Second shell = 8 electrons
Third shell = 8 electrons.

11 electrons




1st shell = 2
2nd shell = 8
3rd shell = 1

Total = 11 electrons



Plum pudding model



Differences to nuclear model

- Ball of positive charge (no protons)
- No nucleus
- No neutrons
- Evenly distributed mass

Rutherford tested the plum pudding model

History of the atom

Scientist	Time	Discovery
John Dalton	Start of the 19 th century	Atoms were first described as solid spheres.
JJ Thomson	1897	Plum pudding model – atom is a ball of + charge with electrons scattered
Ernest Rutherford	1909	Alpha scattering experiment - mass concentrated at the centre, only the nucleus is + charged. Most of the atoms is empty space.
Niels Bohr	Around 1911	Electrons are in shells orbiting the nucleus
James Chadwick	Around 1940	Discovered that there are neutrons in the nucleus.


Rutherford's scattering experiment

alpha particles are positively charged

Fired at gold foil

some alpha particles are deflected/ repelled

most alpha particles passed straight through



What happened?

Conclusions made

Observation	Conclusion
Most of the particles passed straight through	Most of the atom is empty space
Some were deflected to the sides	The particles had passed close by a positive charge
A very small number were repelled straight back	The alpha particles had approached the nucleus straight on. the tiny number told him that the positive charge is in a very small dense core

- Elements **are less** reactive because:
- More electron shells
- Outer shell is further from nucleus and is **more shielded** by the other shells
- The electrostatic force of attraction between free electron and nucleus is **weaker**
- Harder to attract an electron into the outer shell.

C2 – Bonding, structure, and the properties of matter

Formation of Ions

- **Ions** = a charged particle made when atoms lose or gain electrons
- **Positive ion** = atom has lost electrons
- **Negative ion** = atom has gained electrons.

Metals form **positive ions**

Non-metals form negative ions


Group	Ions	Example
1	+1	$\text{Li} \rightarrow \text{Li}^+ + \text{e}^-$
2	+2	$\text{Ca} \rightarrow \text{Ca}^{2+} + 2\text{e}^-$
6	-2	$\text{O} + 2\text{e}^- \rightarrow \text{O}^{2-}$
7	-1	$\text{Br} + \text{e}^- \rightarrow \text{Br}^-$

Lost electrons

Gained electrons

Metallic Bonding


- Happens in **metals only**.
- Positive metal ions surrounded by **sea of delocalised electrons (can move)**.
- Ions tightly packed in rows.
- Strong **electrostatic forces of attraction** between positive ions and negative electrons.




Alloys

- **Alloys** = mixture of two or more metal atoms
- Pure metals are too soft for many uses.

Pure Metal




Alloy



- | | |
|-------------------|-------------------------|
| • Atoms same size | • Different sized atoms |
| • Layers slide | • Layers cannot slide |
| • Softer | • Stronger |

Ionic Bonding


- Between a metal and non-metal.
- Metals give electrons to non-metals so both have a full outer shell.
- **Electrostatic force of attraction** between positive and negative ions.



E.g. Sodium loses one electron to become Na^+ . Chlorine gains one electron to become Cl^- . The two ions attract to form sodium chloride.

Ionic compounds

- Form **giant lattices**, as the attraction between ions acts in all directions



Properties of Ionic Compounds


- **High melting point** – lots of energy needed to overcome electrostatic forces.
- **High boiling point**
- **Cannot conduct electricity as solid** – ions cannot move
- **Conducts electricity when molten or dissolved** – ions are free to move.

Covalent Bonding

- **Covalent bonding** = sharing a pair or pairs of electrons for a full outer shell.
- Between **non-metals only**.

Dot and cross diagrams

- Show the bonding in simple molecules.
- Uses the outer shell of the atoms
- Crosses and dots used to show electrons
- You should be able to draw the following:




Simple Covalent Molecules

- Form when all atoms have full outer shells so bonding stops
- Examples are the molecules shown above.
- Have **low melting and boiling points**
- Due to **weak intermolecular forces**
- Do not conduct electricity

C2 – Bonding, structure, and the properties of matter


Giant Covalent Structure – Diamond

- Each carbon atom **covalently** bonded to **four** others.
- Forms a giant structure
- This makes diamond **strong** → a lot of **energy** needed to break lots of strong covalent bonds.
- **Does not conduct electricity** – has no free electrons.



Giant Covalent Structure – Graphite


- Layers of **carbon** arranged in **hexagons**.
- Each carbon bonded to **three** other carbons.
- Leaves **one delocalised electron** → moves to carry electrical charge **throughout structure**.



- Layers held together by **weak forces**
- Layers can **slide** over each other easily
- Makes graphite **soft/slippery** → good lubricant.
- Has **high melting point** as has many strong covalent bonds.

Silicon Dioxide


- Similar structure to diamond
- Giant covalent structure.
- Lots of **strong covalent bonds**.
- These require lots of **energy** to break.
- High melting and boiling points.



Fullerenes and Nanotubes

- Molecules of carbon shaped into hollow tubes or balls.
- Used to **deliver drugs into body**


Buckminsterfullerene
Formula = C₆₀



- **Carbon nanotubes** = long narrow tubes
- Can conduct electricity


- Can strengthen materials without adding weight.

- Used in electronics and nanotechnology.



Graphene



- Graphene = one layer of graphite.
- Very strong → lots of strong covalent bonds.



- Each carbon bonded to three others.
- One **free delocalised electron** → can move to **carry electrical current** throughout the structure.

Molecular models


- There are different ways to show a molecule other than dot and cross diagrams.



C2 – Bonding, structure, and the properties of matter

States of Matter

- Three states of matter: **solid, liquid & gas.**
- To change state, **energy** must be **transferred.**



- When heated, particles **gain energy.**
- **Attractive forces** between particles begin breaking when melting or boiling points are reached
- **Amount of energy** needed to change state depends on how strong forces are.

Gas


- Randomly arranged.
- Particles **move quickly** – all directions.
- Highest **amount of kinetic energy.**



- Gases **are able to flow** – fill containers
- **Can be compressed** as there is **space between particles**

Solid


- **Regular** pattern (rows and columns)
- Particles **vibrate** in a **fixed position.**
- Particles have **low amount of kinetic energy.**



- Have a **fixed shape** – cannot flow because of strong forces of attraction between particles
- **Cannot be compressed** – particles close together.

Liquid

- Particles **randomly** arranged and touching.
- Particles can **move around.**
- **Greater amount of kinetic energy** than solid



- Liquids **able to flow** – take shape of containers.
- **Cannot be compressed** – particles are close together and cannot be pushed closer

State symbols

- States of matter shown in chemical equations:
- Solid (**s**)
- Liquid (**l**)
- Gas (**g**)
- Aqueous (**aq**)
- **Aqueous** solutions = substance dissolved in water.

Identifying Physical State of Substances

- If the temperature is **lower** than a substance's melting point – substance is **solid.**
- If the temperature is **between** the melting point and boiling point – substance is **liquid.**
- If the temperature is **higher** than the boiling point – substance is a **gas.**

Limitations of Particle Model (HT)

- No chemical bonds are shown.
- Particles shown as solid spheres – not the case, particles are mostly empty space like atoms.
- The diagrams don't show any of the forces between particles
- The diagrams are unable to show the movement of the particles.


C3 – Quantitative Chemistry

Conservation of Mass

- Atoms cannot be created or destroyed during reactions.
- **Mass of reactants = mass of products.**


To show mass is conserved in a reaction:

M_r on the left-side must be same as the right side.



Reacting masses

Use conservation of mass to predict masses:



Both sides need to be equal:
 $10\text{g} - 8\text{g} = 4\text{g}$ of oxygen

Percentage Mass

- Percentage mass of an element in a compound

$$\frac{\text{Mass of the element in compound}}{\text{Total mass of compound}} \times 100$$

Example Question:

Find the percentage mass of oxygen in magnesium oxide (MgO).

A_r of magnesium = 24 A_r of oxygen = 16

M_r of MgO = $24 + 16 = 40$

$$\% \text{ mass} = \frac{A_r}{M_r} = \frac{16}{40} = 0.4 \times 100 = 40\%$$

X 100 to make a % 40% of the mass of MgO is oxygen

Mass Changes

- Mass is always conserved in a reaction.
- Sometimes it may seem like the mass has increased/decreased.
- If a **reactant** is a gas – mass may **increase**.




Oxygen is in the air before it combines with magnesium – you cannot find the mass of oxygen on the balance.

It will look like the mass has increased when it is re-weighed at the end.



- If a **product** is a gas and the gas is able to escape the system – mass will **decrease**.




It will look like the mass has decreased as some of the atoms have been given off as gas and have escaped – so cannot be re-weighed.

Atomic mass (A_r) and Relative Formula Mass (M_r)

- Atomic mass (A_r) is the mass number – ie the mass of one atom
- Relative formula mass (M_r) = all the **relative atomic masses (A_r)** of the atoms in a compound or molecule added up.

Example



The Mole (HT only)

- **Avogadro constant** – 6.02×10^{23}
- One mole contains 6.02×10^{23} atoms or molecules
- The mass, in g, of one mole is the A_r (if an element) or M_r if a compound or molecular element

Iron has a A_r of 56, so 1 mole of iron is 56 g and contains 6.02×10^{23} atoms of iron

Ammonia (NH_3) has an M_r of 17, so 1 mole of ammonia has a mass of 17g. and contains 6.02×10^{23} molecules of ammonia

C3 – Quantitative Chemistry

Concentrations of Solutions

- Concentration = mass of dissolved substance in specific volume (eg dm³)
- More substance dissolved = more concentrated solution

$$\text{Concentration} = \frac{\text{mass}}{\text{volume}}$$

(g/dm³) (g) (dm³)

Can be rearranged to find mass dissolved:

$$\text{mass} = \text{concentration} \times \text{volume}$$

(g) (g/dm³) (dm³)

$$1000\text{cm}^3 = 1\text{dm}^3$$

$$\text{cm}^3 \rightarrow \text{dm}^3 = \text{divide by } 1000.$$

Calculating mass in a given volume

If you have a known volume of a solution of known concentration then you can calculate the mass of dissolved solid.

E.g Calculate the mass of dissolved solid in 25cm³ of a 96g/dm³ solution

96g/dm³ means 96g in every 1000cm³

Do the same to the other side (÷40)

↓
2.4g

↓
25cm³

How do we get from 1000 to 25? (÷40)

Moles and Equations (HT only)

- You can use moles to help you write balanced symbol equations.

Example Question

18.4g of Sodium reacted with 6.4g of oxygen to give 24.8g sodium oxide. Use the masses to write the balanced equation.

Step	Example
Write the equation for the reaction (unbalanced)	Na + O ₂ → Na ₂ O
write down the mass or % given in the question	18.4 + 6.4 → 24.8
Write the mass of one mole of each element or compound	23 32 62 (e.g 18.4 ÷ 23)
Divide the mass given in question by the mass of one mole	0.8 0.2 0.4
Turn the answers into whole number simple ratio	8 2 4 (cancel down) 4 1 2
Put the numbers into the equation	4Na + O ₂ → 2Na ₂ O

Calculating reacting masses (HT)

Example Question

Calculate the mass of calcium needed to make 11.2g Calcium oxide

Step	Calculation
Write the balanced equation	2Ca + O ₂ → 2CaO
Write the masses of each substance	80 + 32 → 112
Write down the given mass in the question.	11.2
Work out the 'scale' factor (ie what did you have to do to the original number to get to the desired mass	÷ 10
Do the same to the other side	8g

Limiting Reactants (HT only)

- If one reactant runs out before the other, then the reaction will stop.
- The reactant that runs out first in a reaction is known as the limiting reactant.

C4 – Chemical Changes


The Reactivity Series

- A more reactive metal will replace a less reactive metal in a compound (**displacement**)

- e.g. potassium + magnesium chloride → potassium chloride + magnesium

Potassium is more reactive than magnesium

Potassium **displaces** magnesium from the compound and takes its place.



Reactions of acids with metals

- Metal + acid → salt + hydrogen

E.g. iron + sulfuric acid → iron sulfate + hydrogen

iron sulfate

salt

To name salt :
1st name Metal
2nd name Acid used

Naming Salts

Acid used	Salt produced
Hydrochloric	Chloride
Sulfuric	Sulfate
Nitric	Nitrate

Extraction of Metals

- Extraction = remove metal from an ore or a compound.

Ore = a rock containing enough metal to make extracting metal worthwhile.

How to extract metals:

Less reactive than carbon – reduction with carbon

Reduction = loss of oxygen

E.g. iron oxide + carbon → iron + carbon dioxide

Oxygen has been removed to extract iron.

Carbon and the oxygen removed from the iron react to make carbon dioxide


More reactive than carbon – electrolysis is used.

- Some metals are found in **native** form (not reacted, so in element form) – usually platinum and gold as **very unreactive**.

Reaction of metals with oxygen

- Metal + oxygen → metal oxide

e.g. magnesium + oxygen → magnesium oxide



Oxidation reaction as metal gained oxygen

- Oxidation = gaining oxygen

- Reduction = losing oxygen


Reaction of metals with water

- Most metals don't react well with water

- Group 1 and group 2 react to form alkalis

- Metal + water → metal hydroxide + hydrogen

e.g. lithium + water → lithium hydroxide + hydrogen



Metal hydroxides are alkaline

Reactions of acids with alkalis


- Acid + alkali → salt + water

neutralisation

Hydrochloric acid + sodium hydroxide → sodium chloride + water

sodium chloride

salt



Reactions of acids with carbonates

- Acid + carbonate → salt + water + carbon dioxide

sulfuric acid + calcium carbonate → calcium chloride + water + carbon dioxide

calcium chloride

salt




C4 – Chemical Changes

Redox Reactions (HT only)

- Redox = reduction and oxidation takes place at same time in a reaction.

- Metal + acid = redox reaction

Example



Ionic equation: $2\text{H}^+ + \text{Ca} \rightarrow \text{Ca}^{2+} + \text{H}_2$ Lost 2 electrons (oxidation)

Half equation 1: $\text{Ca} \rightarrow \text{Ca}^{2+} + 2\text{e}^-$

Half equation 2: $2\text{H}^+ + 2\text{e}^- \rightarrow \text{H}_2$ Gained 2 electrons (reduction)

Strong/Weak Acids (HT only)

Strong acid = completely dissociates in a solution
e.g. $\text{HCl} \rightarrow \text{H}^+ + \text{Cl}^-$

Examples = nitric acid and sulfuric acid

Weak acid = partially dissociates in solution.

e.g. $\text{CH}_3\text{COOH} \rightleftharpoons \text{CH}_3\text{COO}^- + \text{H}^+$
 $\rightleftharpoons$ = reversible reaction

Hasn't fully turned into ions – only partially


Concentration = how much is dissolved in every cm^3

Strong/weak = how well it ionises

As **pH** decreases by 1 unit, **hydrogen ion concentration** of solution increases by factor of 10

Electrolysis


- **Splitting** up a **compound** using electricity.
- Used to extract metals from compounds, purify metals (eg copper)



- Must be **molten** or **aqueous** (dissolved in water) to allow **ions** to **move** to the electrodes


The Process of Electrolysis

Two **electrodes** – made of **inert** material (doesn't react)



pH Scale


- Shows how acidic or alkaline solution is.
- pH 1-6 = acid
- pH 7 = neutral
- pH 8-14 = alkali



In aqueous solutions:


Acids – produce H^+ ions
Alkalis – produce OH^- ions

In neutralisation reactions:



Measuring pH of a solution

- Can use **universal indicator**
- Gives the solution a colour
- Can compare colour to the pH scale



Disadvantages of method

- Colour is **subjective** – different people may see different colours
- Doesn't give an exact pH number (could use **pH probe** to make more **accurate**).

C4 – Chemical Changes – Required Practical – Preparation of soluble salts

Aim

Prepare a pure, dry sample of a soluble salt from an insoluble **oxide or carbonate**.


Equipment

- Beaker
- Measuring cylinder
- Bunsen burner and safety mat
- Filter funnel and filter paper
- Named acid (e.g. hydrochloric acid)
- Metal oxide or carbonate.
- Spatula
- Glass stirring rod

Change method
depending on reactants in
the question.

Method (example copper oxide and sulfuric acid to make copper sulfate)

1. Using measuring cylinder – 20cm³ **sulfuric acid** → beaker
2. Warm the acid gently (not boiling)
3. Using spatula add **copper oxide** to the acid and stir
4. Keep adding until no more oxide will dissolve (excess).
5. Using a filter funnel and filter paper – filter excess copper oxide.
6. Evaporate some of the filtrate using a water bath.
7. Pour remaining filtrate into an evaporating basin – leave overnight to evaporate water
8. Pat the crystals dry.



Common questions

Q1) Why do you heat the acid before adding the oxide?

A1) To speed up the reaction (particles have more energy to react).

Q2) Why is the oxide added in excess?

A2) To make sure that all the acid has been neutralised.

Q3) Why is the solution filtered?

A3) Remove any unreacted, excess solid.

Q4) Why is the solution left overnight in a warm, dry place?

A4) To evaporate excess water, to form crystals (crystallise).

Q5) Name 2 safety precautions you should take during this practical.

A5) Safety goggles and allow equipment to cool before putting away

C4 – Chemical Changes


Electrolysis of Molten Ionic Compounds

Molten = melted so ions can move.

- Metal = produced at **anode**
- Non-metal = produced at **cathode**

Example: Lead Bromide - PbBr_2

Lead forms at the cathode
Bromine gas is formed at anode



Using Electrolysis to Extract Metals

- Used if metal is **too reactive** to be extracted by reduction with carbon.
- Requires **large amount of energy** to melt the compound and produce electrical current. (**expensive**)


Example: Aluminium Oxide

- **Cryolite** is added – reduces the melting point (less energy needed – less expensive)
- **Carbon** used as positive electrode – needs to be replaced constantly as **oxygen** will react with it to produce CO_2 – it will degrade.

Electrolysis of Aqueous Solutions

- Compound is dissolved in water so ions can move.

When aqueous – H^+ and OH^- (from H_2O) are also present along with the two ions from the compound.



- Only **one** ion is discharged at each electrode.

Anode – Non-metal or oxygen

Cathode – Metal or hydrogen

Rules

+ ANODE Attracts – ions ('Anions')	- CATHODE Attracts + ions ('Cations')
If – ions are group 7 i.e. chloride Cl^- bromide Br^- iodide I^- Then the groups 7 element is produced as a gas	If + ions (metals) are MORE REACTIVE than hydrogen K, Na, Ca, Mg, Zn, Fe Then HYDROGEN is produced
If – ions are NOT Group 7 Eg sulphate SO_4^{2-} nitrate NO_3^- carbonate CO_3^{2-} OXYGEN is produced.	If + ions (metals) are LESS REACTIVE than hydrogen Cu, Ag, Au Then the METAL is produced

Examples

Solution	Product at cathode	Product at anode
Potassium chloride	Hydrogen – because K is more reactive than H	Chlorine – as it is a halogen
Copper sulfate	Copper – as copper is less reactive than H	Oxygen – as there is no halogen

Half-Equations at Electrodes (HT only)

During electrolysis:


Cathode – positive ions **gain** electrons (**reduction**)

Anode – negative ions **lose** electrons (**oxidation**)

- Ions become **discharged** (lose their charge) at the electrodes to form the atoms again.


- Reactions at electrodes can be represented by half equations.

Examples




Gained 2 electrons (reduction)

molecules of hydrogen gas produced




molecules of oxygen produced

Lost electrons (oxidation)



Gained electrons (reduction)

Copper atoms are formed at the cathode



chlorine molecules are formed

Lost electrons (oxidation)

C4 – Chemical Changes – Required Practical – Electrolysis of Aqueous Solutions

Aim

To investigate the electrolysis of an aqueous solution using inert (unreactive) **electrodes**.


Equipment

- Beaker
- Two test tubes (or measuring cylinders)
- Graphite electrodes
- Two splints
- Aqueous solution
- DC powerpack

Change method depending on the question.

Method (example copper sulfate solution.)

1. Pour some copper sulfate solution into a beaker.
2. Place two graphite rods into the copper sulfate solution. Attach one electrode to the negative terminal of a dc supply, and the other electrode to the positive terminal.
3. Completely fill two small test tubes with copper sulfate solution and position a test tube over each electrode as shown in the diagram. **(use measuring cylinders if measuring volume of gas produced)**
4. Turn on the power supply and observe what happens at each electrode.
5. Test any gas produced with a glowing splint and a burning splint.
6. Record observations and the results of your tests.



Common questions

Q1) How do you test for hydrogen gas?

A1) Lit splint will make a squeaky pop.

Q2) How do you test for oxygen gas?

A2) Glowing splint – will relight.

Q3) Explain why copper is produced at the cathode.

A3) Copper ions are **positive**, so are attracted to the negative electrode (opposites attract). Copper is less reactive than hydrogen so is discharged. The copper ions **gain electrons** and are **reduced** to form **copper atoms**.

Q4) Why do hydrogen ions move to the cathode?

A4) Hydrogen ions are **positive** so move to the negative electrode as **opposites attract**.

Q5) Why are measuring cylinders better to collect the gas?

A5) Because they are more accurate when measuring the volume of gas produced.

C5 – Energy Changes

Exothermic Reactions

- Energy transferred to the surroundings
- Temperature of the reaction mixture **increases**
- This energy is transferred **to** the surroundings

Examples include:

- Hand warmers
- Combustion reactions
- Respiration
- Neutralisation reactions
- Self-heating cans.



Exothermic

Endothermic Reactions

- Energy absorbed from the surroundings
- Temperature of reaction mixture often **decreases**
- Energy is transferred **from** the surroundings

Examples include:

- Ice packs (injuries)
- Reaction of citric acid and sodium hydrogen carbonate
- Thermal decomposition of calcium carbonate



Endothermic

Energy change of reactions (HT)

During a reaction:

- Energy is **absorbed** in order to **break** bonds in the reactants 
- Energy is **released** when bonds are **made** in the products. 


Bond energy = the amount of energy that is released when a bond is made or that is needed to break a bond

Calculating energy changes (HT)

Overall energy change = difference between energy needed to break bonds and the energy **released** when bonds formed.

To calculate energy change :

Energy change = bonds broken – bonds formed



Bond	Bond Energy / kJ mol^{-1}
F—F	158
H—H	436
H—F	568


Bonds broken = $436 + 158$ 593	Bonds formed 2×568 1136
--	--

**Overall energy change = $593 - 1136$
= -543 kJ/mol Exothermic**

More energy is released in bond making than is required for bond breaking.

Reaction Profiles – Exothermic


- Energy level diagrams show **difference in energy** between reactants and products.
- Exothermic = Energy of products is **lower than** reactants (energy is released)
- **Activation Energy** = minimum amount of energy needed to start the reaction.
- **Energy change** = the difference in energy between reactants and products.



You may need to draw and label this in the exam!

Reaction Profiles – Endothermic

- Energy level diagrams show **difference in energy** between reactants and products.
- Endothermic = Energy of products is **higher than** reactants (energy is absorbed)
- **Activation Energy** = minimum amount of energy needed to start the reaction
- **Energy change** = the difference in energy between reactants and products.



You may need to draw and label this in the exam!

C5 – Energy Changes – Required Practical – Temperature Changes

Hypothesis

The energy change in the reaction between acid and alkali depends on the volume of alkali added.

Equipment

- Polystyrene cup and lid
- Thermometer
- 250cm³ beaker
- Measuring cylinder
- Liquid reactants



Method (example for hydrochloric acid and sodium hydroxide)

1. Using measuring cylinder to measure 30cm³ hydrochloric acid and put in polystyrene cup
2. Stand cup inside beaker to make stable.
3. Use a thermometer to measure the temperature of acid and record.
4. Using measuring cylinder – 5cm³ sodium hydroxide → polystyrene cup
5. Fit the lid and gently stir with thermometer through hole.
6. When reading stops on thermometer, record temperature in table.
7. Repeat, each time adding 5cm³ more sodium hydroxide up to a maximum of 40cm³.
8. Calculate the temperature change on each attempt.
9. Repeat the experiment 3 times and calculate a mean temperature change for each volume of sodium hydroxide.

Variables

Independent – Volume of sodium hydroxide

Dependent – Temperature change

Control – Volume of hydrochloric acid, concentration of acid, concentration of sodium hydroxide

Common questions

Q1) Why do you use a polystyrene cup and lid?

A1) Because polystyrene cups are insulators, which reduces heat loss in the experiment, making the results more accurate.

Q2) Why should you calculate the temperature change, instead of just using the final temperature?

A2) Because the initial (starting) temperature of the acid may have been different.

Q3) Why is it important to stir the mixture?

A3) To make sure all of the reactants have reacted and to get a uniform temperature.

Q4) Why is the experiment conducted 3 times?

A4) So that anomalies can be seen and removed and a mean calculated

Energy changes could also be investigated using:

1. Changing the **mass of metal** added to acid and measuring the **temperature increase**
2. Changing the **type of metal** added to acid and measuring the **temperature increase**
3. Dissolving different **masses of potassium nitrate** into water and observing the **temperature decrease**.